

Experimental Evidence of “Granulence”

G. Combe*, V. Richefeu*, G. Viggiani*, S.A. Hall†, A. Tengattini* and
A.P.F. Atman**

*UJF-Grenoble 1, Grenoble-INP, CNRS UMR 5521, 3SR Lab., B.P. 53, 38041 Grenoble Cedex 09, France.

†Division of Solid Mechanics, Lund University, Lund, Sweden and European Spallation Source AB, Lund, Sweden

**Departamento de Física e Matemática and National Institute of Science and Technology for Complex Systems, Centro Federal de Educação Tecnológica de Minas Gerais, CEFET-MG, Av. Amazonas 7675, 30510-000, Belo Horizonte-MG, Brazil

Abstract. We present an experimental study of velocity fluctuations in quasistatic flow of a 2D granular material deformed in a shear apparatus named $1\gamma 2\epsilon$ [1]. Radjaï and Roux [2] revealed systematic similarities between velocity fluctuations observed in discrete element simulations of quasistatic flow of granular material and turbulent flows in fluids. The character of these velocity fluctuations – named *granulence* by [2] – manifests as a non-Gaussian broadening of the probability density function of the fluctuations as the length of the analyzed shear-window is decreased, and exhibits some space and time scaling. The experiments presented are simple shear tests on granular samples composed of about 2000 wooden rods. The kinematics of the rod centers was followed by means of 2D Particle Image Tracking (PIT) technique applied to a sequence of 24 Mpixels digital pictures acquired throughout the duration of the loading at a frequency of 0.08 image/s. This analysis confirms the existence of granulence features in a real experimental test, which is comparable to that previously observed in numerical simulations of [2]. The experimental results obtained open up a new avenue for further studies on fluctuations in granular materials.

Keywords: Granular materials, Quasi-static shear, Kinematic fluctuations, Spatial Correlation, Structured Patterns

PACS: 45.70.-n, 45.70.Qj, 45.50.Dd, 46.65.+g, 05.40.-a

INTRODUCTION

In a granular material, a macroscopically homogeneous deformation does not correspond to a homogeneous field of displacement gradient when looking at the individual grains. Due to geometrical constraints at the grain scale (mutual exclusion of grain volumes), grains are not able to displace as continuum mechanics dictates they should. This can be pictured by imagining an individual in a crowd of people who all wish to go to the same place: although the long-term displacement of each individual is equal to the displacement of the crowd, the steps of each individual are erratic. Classical continuum approaches disregard this feature, which is apparent only at “small” strains (*i.e.*, for displacements that are small with respect to the size of rigid grains, or the size of contact indentation for deformable grains). However, the deviation of a grain’s displacement from the value dictated by the continuum field (referred to as *fluctuation*) is likely to hold valuable information about the characteristic length(s) involved in grains’ rearrangement, which is the principal mechanism of irreversible deformation in granular materials. Displacement fluctuations, often spatially organized in the form of vortices, have been observed in (quasistatic) experiments only by [3]. Several numerical studies using Discrete Elements (DEM) report similar observations (e.g., [4, 5, 6, 7, 8]). [2] investigated

such fluctuations by making use of statistical analysis that is typically used in fluid dynamics. The main finding of [2] was that displacement fluctuations in granular materials show scaling features with striking analogies to fluid turbulence. Inspired by the approach of [2], the present study analyzes fluctuations measured from experiments on a 2D analogue granular material subjected to (quasi-static) shear.

SHEAR TEST

Shear tests have been performed in the $1\gamma 2\epsilon$ apparatus, which is essentially a plane stress version of the directional shear cell developed for testing soils (see [1, 9, 10] for details). Results from only one such test are discussed herein, although the results are consistent throughout the entire campaign. In this experiment, an assembly of 2000 2D “grains” (wooden cylinders 6cm long) having four different diameters (8, 12, 16 and 20 mm), was slowly deformed in simple shear at constant vertical stress $\sigma_n = 50$ kPa, Fig. 1. Note that the contact properties among the (wooden) grains compare with the imposed in DEM simulations properties used by [2]. The vertical sides of the enclosing frame are tilted up to $\gamma = 0.26$ (15°) while the length of the horizontal sides is kept constant. The shear strain rate is $8.2 \times 10^{-5} s^{-1}$, which is small enough to en-




FIGURE 1. Sketch of a shear test in the $1\gamma 2\varepsilon$ apparatus.

sure a quasi-static regime according to the inertial number criterion suggested by [6]. A video of the test can be found at <http://youtu.be/B4Dfesn5vhs>.

During the test, the normal (σ_n) and tangential (σ_t) stresses are measured at the boundary of the sample (shown in [11]). The global stress-strain response is typical of a dense 2D granular material, with a peak friction angle of about 26° at $\gamma \simeq 0.06$ and a dilatant behaviour throughout. A digital camera was used to acquire 24.5 Mpixels images every 5 seconds (*i.e.*, a shear strain increase of $\Delta\gamma \simeq 4 \times 10^{-4}$, called strain window in the following) throughout the test.

The displacements of all the grains are measured by means of a software named TRACKER specially developed to process the digital images and measure (with sub-pixel resolution) the in-plane displacement and rotation of each individual grain from one image to another. This software uses a discrete grain-scale version of digital image correlation technique hereafter named PIT (Particle Image Tracking) that is well suited for tracking rigid particles. The principle of the PIT technique is detailed in [11] and [12]. From the 24.5 Mpixels images taken during the shear test, particles' displacements are assessed with an error that is less than 0.1 pixel [11]. As an example, Fig. 2 shows the displacements of all grains measured by PIT from 0 to 0.12 shear strain γ .

FLUCTUATIONS OF DISPLACEMENT

To assess the fluctuations of displacement (angular rotations of grains are also assessed, but not discussed in this paper) in the course of deformation, we consider two possible displacements of each grain during a strain window $\Delta\gamma$ (always positive). The first is the actual displacement vector $\delta\mathbf{r}(\gamma, \Delta\gamma)$, which depends both on the size of the strain window $\Delta\gamma$ and the level of shear strain γ at the beginning of the strain window. The second displacement vector $\delta\mathbf{r}^*(\gamma, \Delta\gamma)$ is the displacement dictated by a homogeneous (*affine*) continuum strain field, *i.e.*, the displacement that the grain's center would have if it moved




FIGURE 2. Particles displacement (maximum displacement = 46mm) in a sheared 2D granular assembly made of 2000 wooden rods. The granular packing is enclosed by a rigid frame initially rectangular ($0.56\text{m} \times 0.47\text{m}$). A speckle of black and white points is painted on each cylinder to allow the measurement of particle kinematics by means of the PIT technique.




FIGURE 3. Displacement fluctuations $u(\Delta\gamma = 0.1, \gamma = 0)$ of grains in a sheared 2D granular assembly. The corresponding grains' displacement are shown in Fig. 2. The maximum value of u is 18mm .

as a material point in a continuum. The fluctuation of the displacement is defined as the difference between these two displacement vector:

$$\mathbf{u}(\gamma, \Delta\gamma) = \delta\mathbf{r}(\gamma, \Delta\gamma) - \delta\mathbf{r}^*(\gamma, \Delta\gamma). \quad (1)$$

Figure 3 shows a map of the fluctuations u associated with the displacement field of Fig. 2. Displacement fluctuations can be conveniently normalized by dividing the vector $\mathbf{u}(\gamma, \Delta\gamma)$ by the product $\Delta\gamma \langle d \rangle$ (where $\langle d \rangle$ is the mean diameter of the grains), which can be interpreted as the average displacement of the grains in the strain window $\Delta\gamma$. This normalized fluctuation,

$$\mathbf{V}(\gamma, \Delta\gamma) = \frac{\mathbf{u}(\gamma, \Delta\gamma)}{\Delta\gamma \langle d \rangle}, \quad (2)$$

can also be interpreted as a local (microscopic) strain fluctuation, which is in turn divided by the size of the global (macroscopic) strain window $\Delta\gamma$.

The mean displacement fluctuation $\langle u \rangle$ increases monotonically throughout the test, from 0.066 mm (for $\Delta\gamma = 10^{-3}$) to 6 mm (for $\Delta\gamma = 0.25$), this final value correspond to about 10% of the average displacement of the grains between the beginning and the end of the shear test. Note that the smallest displacement fluctuation is well above the accuracy of TRACKER. As far as the mean magnitude of normalized fluctuations $\langle V \rangle$ is concerned, a decrease of $\langle V \rangle$ from 3 to 2 is observed; this implies that the average fluctuation of local shear strain $\langle u \rangle / \langle d \rangle$ is two to three times larger than the global strain window $\Delta\gamma$.

PDF OF THE FLUCTUATIONS

Figure 4 shows the probability density function (pdf) of normalized fluctuations \mathbf{V} along the x direction for two different strain windows $\Delta\gamma$. Whatever the strain window, a good candidate for the *pdf* is

$$F(V_x, q) = a [1 + b(1 - q)V_x^2]^{-\frac{1}{1-q}} \quad (3)$$

This function, named *q-gaussian*, comes from the generalized thermodynamics theory, which is derived from the principle of non-extensive entropy introduced by [13]. The parameter q in Eq. 3 quantifies the degree of non-extensivity of the entropy of the system. In the limit of $q \rightarrow 1$, the function tends to a *Gaussian* probability density function.

Statistical analysis of the fluctuations shows that whatever the value of $\Delta\gamma$, their spatial average is zero, which is expected for a global homogeneous deformation. A key observation is that the pdf exhibits a wider range of fluctuations with decreasing $\Delta\gamma$, which corresponds to a variation of q from 1.14 to 1.59. In [11], a *Kurtosis* analysis of V_x values was seen to tend to zero asymptotically with increasing size of $\Delta\gamma$. A zero value of kurtosis might indicate that the pdf tends to become Gaussian for $\Delta\gamma \rightarrow \infty$, which is confirmed by the decrease of q when $\Delta\gamma$ increases. Very similar *Kurtosis* features and trends have been observed (on DEM simulations) by [2], who noted that they have striking similarities with turbulent flow of fluids (although no dynamics are associated with these fluctuations).




FIGURE 4. Probability density function (pdf) of normalized fluctuations V_x projected on the horizontal direction for two different strain windows, $\Delta\gamma = 2 \times 10^{-3}$ and $\Delta\gamma = 10^{-1}$.

SPATIAL CORRELATIONS OF THE FLUCTUATIONS

To understand the fluctuations, the analysis of their statistics must be supplemented by the study of their spatial distribution.

Spatial correlation of \mathbf{V} can be seen, *e.g.*, the “loop patterns” in Fig. 3. Figure 5 shows three normalized fluctuation maps for three different values of γ , for two different sizes of the strain window $\Delta\gamma$. On the left, the two fluctuation maps computed for $\Delta\gamma = 2 \times 10^{-3}$ at two different values of γ show that both long-range correlation (blue curve, 40 grain diameter of correlation length) and short-range correlation (red curve, 10 diameter of correlation length) are both observed. This is consistent with *Tsallis* theory, which states that when *q*-Gaussian distributions are observed with $q > 1$ (here $q = 1.59$), then long-range spatial correlations emerge. On the right of Fig. 5, the strain windows used to compute the normalized fluctuation is bigger ($\Delta\gamma = 1.7 \times 10^{-1}$); the statistical distribution of V_x is similar to the one shown in Fig. 4 for $\Delta\gamma = 10^{-1}$. The q value of the *q*-Gaussian fit is smaller than for $\Delta\gamma = 2 \times 10^{-3}$ ($q = 1.14$), but still greater than 1 (perfect *Gaussian* distribution). A strong spatial correlation of V_x is found for distances less than $\Delta = 10$ particle diameters. This length roughly corresponds to the minimum radius of the fluctuation loops. A pseudo-period of about $\Delta = 20$ diameters is observed, which is another signature of the loops. This pseudo-periodicity might indicate the existence of a cascade of loop-sizes rather than a unique size – the smallest size being about 10 diameters, and the largest in the order of the sample size. This interpretation was confirmed by [11] with a Fourier transform of the fluctuation signal as a function of space, as originally suggested by [2].




FIGURE 5. Maps of the normalised fluctuations for two different values of $\Delta\gamma$. The curves show the spatial auto-correlogram for the x component of \mathbf{V} . Δ is the correlation length expressed in terms of grain diameters.

CONCLUSIONS

A detailed study of fluctuations of displacements in a granular material subjected to quasistatic shear has revealed that the fluctuations are organized in space and show clear vortex-like patterns, reminiscent of turbulence in fluid dynamics. While these fluctuation loops have been often observed in DEM numerical simulations, on the experimental front they have only been reported (to the authors' best knowledge) in the study by [3]. The present study (results of which have already been presented in [11]) confirms these early findings and brings into the picture the important idea that fluctuation loops have a characteristic minimum radius – equal to 10 mean particle diameters for the material tested. Displacement fluctuations in granular materials are a direct manifestation of grain rearrangement, therefore they can be thought of as the basic mechanism of irreversible deformation. The statistical analysis of these fluctuation was described by q -Gaussian probability distribution, which is a signature of long-range correlation for $q > 1$. The link between these fluctuations (and their spatial organization) and the deformation of granular materials at the macro scale will be investigated in future work.

ACKNOWLEDGMENTS

The authors are indebted to C. Tsallis by very fruitful discussions. G. Combe thanks to the Centro Federal de Educação Tecnológica de Minas Gerais (Brazil) for financial support. A.P.F. Atman thanks to CNPq and FAPEMIG Brazilian agencies for financial support.

REFERENCES

1. F. Calvetti, G. Combe, and J. Lanier, *Mechanics of Cohesive Frictional Materials* **2**, 121–163 (1997).
2. F. Radjai, and S. Roux, *Phys. Rev. Lett.* **89**, 064302 (2002).
3. A. Misra, and H. Jiang, *Computers and Geotechnics* **20**, 267–285 (1997).
4. J. R. Williams, and N. Rege, *Mechanics of Cohesive-frictional Materials* **2**, 223–236 (1997).
5. M. R. Kuhn, *Mechanics of Materials* **31**, 407–429 (1999).
6. G. Combe, and J.-N. Roux, “Discrete numerical simulation, quasistatic deformation and the origins of strain in granular materials,” in *3rd International Symposium on Deformation Characteristics of Geomaterials*, edited by di Benedetto et al., Lyon, 2003, pp. 1071–1078.
7. A. Tordesillas, M. Muthuswamy, and S. Walsh, *Journal of Engineering Mechanics* **134**, 1095–1113 (2008).
8. A. L. Rechenmacher, S. Abedi, O. Chupin, and A. D. Orlando, *Acta Geotechnica* **6**, 205–217 (2011).
9. H. Joer, J. Lanier, J. Desrues, and E. Flavigny, *Geotechnical Testing Journal* **15**, 129–137 (1992).
10. E.-M. Charalampidou, G. Combe, G. Viggiani, and J. Lanier, “Mechanical behavior of mixtures of circular and rectangular 2D particles,” in *Powders and Grains 2009*, edited by M. Nakagawa, and S. Luding, AIP, Golden, USA, 2009, pp. 821–824.
11. V. Richefeu, G. Combe, and G. Viggiani, *Géotechnique Letters* **2**, 113–118 (2012).
12. G. Combe, and V. Richefeu, “TRACKER: a Particle Image Tracking (PIT) technique dedicated to nonsmooth motions involved in granular packings,” in *Powders and Grains 2013*, AIP, Sydney, Australia, 2013.
13. C. Tsallis, *Journal of Statistical Physics* **52**, 479–487 (1988).